



Taxonomic paper

The millipede *Typhloglomeris caucasica* Golovatch, 1975 found epigeically (Diplopoda, Glomerida, Glomeridellidae)

Sergei Golovatch[†], Yuri A. Chumachenko[‡]

[†] Russian Academy of Sciences, Moscow, Russia

[‡] Caucasian State Biosphere Nature Reserve, Research Department, Sovetskaya Street 187, Maikop, Republic of Adygeya, Russia

Corresponding author: Sergei Golovatch (sgolovatch@yandex.ru)

Academic editor: Robert Mesibov

Received: 02 Aug 2013 | Accepted: 28 Aug 2013 | Published: 16 Sep 2013

Citation: Golovatch S, Chumachenko Y (2013) The millipede *Typhloglomeris caucasica* Golovatch, 1975 found epigeically (Diplopoda, Glomerida, Glomeridellidae). Biodiversity Data Journal 1: e981. doi: [10.3897/BDJ.1.e981](https://doi.org/10.3897/BDJ.1.e981)

Abstract

The millipede *Typhloglomeris caucasica* Golovatch, hitherto considered as a troglobite confined to several caves near Sochi, western Caucasus, Russia, is recorded epigeically in the same region, and is therefore a troglophile.

Keywords

Diplopod, *Typhloglomeris caucasica*, Caucasus, Sochi, cave, forest litter, new record

Introduction

The small diplopod family Glomeridellidae contains only two valid genera: *Glomeridella* Brölemann, 1913, with 7-8 species ranging from France in the west, through the eastern Alps, to the Balkans in the east, and *Typhloglomeris* Verhoeff, 1898, with about 15 species, a few of which are presumed troglobites, from the Balkans, Caucasus, northwestern Iran, and Asia Minor (Golovatch 2003, Makarov et al. 2003). All *Glomeridella* and most

Typhloglomeris species show distinctive colour patterns and are known to be epigean, but even among the few colourless representatives not all are troglobites. Thus, *T. fiumarana* Verhoeff, 1899 has been regarded as possibly a geobiont (Golovatch 1989, Golovatch 2003) because it was found in Croatia under stones, not in a cave. Similarly, *T. alba* (Golovatch, 1989), in which only the ocelli are pigmented, whereas the rest of the body is pallid, is clearly epigean in northwestern Anatolia, Turkey (Golovatch 1989). Only the following fully unpigmented congeners have hitherto been referred to as troglobites (Golovatch 1975, Golovatch 1989, Golovatch 2003, Makarov et al. 2003): *T. coeca* Verhoeff, 1898, from Croatia and Montenegro, *T. varunae* Makarov, Lučić, Tomić & Karaman, 2003, from Macedonia, *T. seuti* Makarov, Lučić, Tomić & Karaman, 2003, from Montenegro, and *T. caucasica* Golovatch, 1975, from near Sochi, western Caucasus, Russia. Furthermore, based on several morphological characters, Makarov et al. 2003 treat *T. caucasica* as representing a species group of its own, the *caucasica*-group.

Taxon treatment

Typhloglomeris caucasica Golovatch, 1975

Material

- a. kingdom: Animalia; phylum: Arthropoda; class: Diplopoda; order: Glomerida; family: Glomeridellidae; genus: *Typhloglomeris*; specificEpithet: *caucasica*; scientificNameAuthorship: Golovatch, 1975; country: Russia; stateProvince: Sochi; verbatimLocality: Khosta, Caucasian Biosphere Nature Reserve; samplingProtocol: pitfall trapping; eventDate: 14 July - 10 August 2006; habitat: *Taxus* & *Buxus* relict forest; individualCount: 1; sex: male; recordedBy: Y. A. Chumachenko; institutionCode: Zoological Museum, Moscow State University, Russia

Taxon discussion

The recent discovery of *T. caucasica* in a pitfall trap in forest near Sochi indicates that the ecological status of this species is a troglophile, not a troglobite, when it occurs in caves in the same area. This record emphasizes the need to carefully collect near and outside caves when assessing the degree of cavernicolity of endogean animals.

Acknowledgements

This work has been supported in part through a grant released to the first author by the Russian Ministry of Education and Science.

References

- Golovatch S (1975) Two genera of Oniscomorpha (Diplopoda) new to the USSR fauna, found in Transcaucasia, and their zoogeographical relations. *Zoologicheskyy zhurnal* 54 (10): 1566-1571. [In Russian, English summary].
- Golovatch S (1989) Diplopoda of the Caucasus, 1. Glomeridellidae, with contributions to the fauna of Turkey. *Senckenbergiana biologica* 69: 405-419.
- Golovatch S (2003) Two new species of Glomeridellidae (Diplopoda: Glomerida) from the Middle East. *Arthropoda Selecta* 11 (4): 255-258.
- Makarov S, Lučić L, Tomić V, Karaman I (2003) Two new glomeridellids (Glomeridellidae, Diplopoda) from Montenegro and Macedonia. *Periodicum biologorum* 105 (4): 473-477.